

Index

A

abstracts

emulsions

- The Influence of Emollients on Skin Penetration from Emulsions, 91
- The Influence of Emulsifiers on the Skin Penetration from Emulsions, 125–126
- The Influence of Emulsion Droplet Size on Cosmetic Delivery, 171
- Liquid Crystalline Emulsions as Cosmetic Delivery Systems, 139–140

encapsulation

- Encapsulation and Other Topical Delivery Systems, 337
- Micro- and Nano-encapsulation of Water- and Oil-soluble Actives for Cosmetic and Pharmaceutical Applications, 313–314
- Microencapsulation and Cosmetics, 251

follicular delivery, Significance of Follicular Delivery and Ways to Measure It, 441

iontophoresis

- Principles of Iontophoresis and its Applications in Cosmetics, 381
- Topical Delivery of Cosmetic Ingredients by Sonophoresis and Iontophoresis, 365

liposomes, Flexible Liposomes for Topical Applications in Cosmetics, 269

percutaneous absorption, Methods for Studying Percutaneous Absorption, 30

Pheroids, The Design of a Skin-Friendly Carrier for Cosmetic Compounds Using Pheroid Technology, 283

rinse-off product delivery, Delivery from Rinse-off Products: Mission Impossible or a Reality for Cosmetic Science, 403

skin and cosmetic delivery, Legal Perspective on Cosmetic and Drug Skin Delivery Systems, 461–462

skin delivery

- Clinical Relevance of Topical Delivery Systems for Active Cosmetic Ingredients, 487
 - Skin Electroporation, 391
- The Influence of Formulation Type on Skin Delivery, 61–62
- Marketing Skin Delivery Systems: Despite So Many Questions, Here Are Some Answers, 477
- Measuring Skin Delivery, 23
- Skin Delivery: What It Is and Why We Need It, 1
- Vesicles as Skin Delivery Vehicles, 205–206

SLN® and SLC®, Challenging Cosmetics—Solid Lipid Nanoparticles (SLN®) and Nanostructured Lipid Carriers (NLC®), 227–229

- sonophoresis, Topical Delivery of Cosmetic Ingredients by Sonophoresis and Iontophoresis, 365
- spectroscopy
- Acoustic Attenuation Spectroscopy: A New Technique to Characterize the Stability and Structure of Semisolid Topical Delivery Systems for Cosmetic and Pharmaceutical Applications, 185–186
 - The Application of Infrared Spectroscopic Imaging to Skin Delivery: Visualizing Molecular Localization in Formulation and in Skin, 40
- stratum corneum
- Delivering Chemically Labile Molecules into the Stratum Corneum: An Example of Stabilizing Linoleic Acid with α -Cyclodextrin, 353
 - The Effect of Changing the Polarity of the Stratum Corneum on the Dermal and Transdermal Delivery of Cosmetically Active Ingredients, 109–110
- transfollicular transport, Novel Ways of Measuring the Transfollicular Transport of Active Ingredients across the Skin, 425–426
- acoustic attenuation spectroscopy, 186–202
- animal skin *versus* human skin, 26–27

B

- biodegradable SLNs (solid-lipid nanoparticles), pharmaceutical and medical uses, 331–334

C

- CAID (cosmetic active ingredient systems). *See also* TDS
- clinical relevance, 488–490
 - percutaneous enhancers, 490–495
- capsules. *See also* encapsulation
- encapsulation (cosmetic), 251–252
 - microencapsulation (cosmetic), 252–253
 - producing, 253–265
- COLIPA (European Cosmetic Toiletry and Perfumery Association), skin penetration studies, 30, 32, 36
- corneocytes, 3
- cosmetic active ingredient systems. *See* CAID
- cosmetic delivery, 1–2
- definition, 2–3
 - dermal, 4, 12
 - delivery guidelines, 15–16
 - delivery measurement, 16–18
 - epidermal, 4
 - legal perspective, 462–464

- antioxidants, 467
- claims, 471–472
- cosmeceuticals, 465–466
- nanotechnology, 472–473
- natural/organic products, 466
- safety testing, 470–471
- skin protectants, 467–468
- test methods, 464–465
- relevance to skin delivery, 10
- 4Rs
 - correct period of time, 6
 - right chemical, 4–5
 - right concentration, 5–6
 - right site of action, 3–4
- skin surface, 11
- stratum corneum, 11–12
- transdermal, 4
 - avoiding, 10
 - measuring, 16–18
- transfollicular, 4
- viable epidermis, 12
- why needed, 2
- cyclodextrins, 358–363

D

- dermaportation, 400
- dermis, 3

E

- eccrine glands, 3
- ECVAM (European Centre for the Validation of Alternative Methods), skin penetration studies, 30
- electroporation of skin
 - combined with other methods, 398
 - of cosmetics, 399
 - dermaportation, 400
 - of macromolecules, 396–397

- patient tolerance, 399
 - of small- to moderately-sized molecules, 394–396
 - ways to enhance, 392–394
- emollient influence, on skin penetration from emulsions, 91–92
 - applied active ingredient doses, 103–104
 - “Formulating for Efficacy,” 98–106
 - skin delivery theory, 92–98
- emulsifiers, influence on skin penetration from emulsions, 126–138
- emulsions
 - droplet size influence on cosmetic delivery, 172–180
 - emollient influence on skin penetration from, 91–92
 - “Formulating for Efficacy,” 98–106
 - skin delivery theory, 92–98
 - emulsifier influence on skin penetration from, 126–138
 - liquid crystalline emulsions as cosmetic delivery systems, 140, 142–155
 - clear gel-like emulsion crystals, 160–164
 - lemellar liquid gel, 164–168
 - molecular assemblies, 140–141
 - multilamellar emulsion of pseudo-stratum corneum lipids, 155–160
- encapsulation, 251–252. *See also* capsules
 - microencapsulation, 252–253, 314–315, 338–341
 - chemical production, 316–317
 - physical production, 317
 - polymeric entrapments, 341–350
 - nanoencapsulation
 - liposomes, 318
 - solid-lipid nanoparticles, 319–331
 - solid-lipid nanoparticles, biodegradable, 331–334
- epidermal delivery, 4
- epidermis, 3
- Ethosomes and Transfersomes, 278–281. *See also* liposomes
- European Centre for the Validation of Alternative Methods. *See* ECVAM
- European Commission Health & Consumer Protection Directorate,
General Guidance Document on Dermal Absorption, 30
- European Cosmetic Toiletry and Perfumery Association. *See* COLIPA

F

- FDA/AAPS, skin penetration studies, 30, 35
- follicular delivery, 441–442. *See also* transfollicular delivery
 - biopsy and measurement of follicular contents, 449–451
 - drug targets, 442–444
 - measuring, 445
 - permeation in vitro skin models, 445–448
 - particle size effects, 453–454
 - physical delivery, 455
 - sebum secretion effects, 454
 - structure and follicle distribution, 444–445
 - in vivo models
 - animal, 451–452
 - human, 453
- “Formulating for Efficacy,” 98–106

H

- hair follicles, skin penetration routes. *See* transfollicular delivery
- human skin *versus* animal skin, 26–27

I

- infinite *versus* finite dose techniques, 33–34
- infrared spectroscopic imaging. *See* IR
- intercellular transepidermal penetration routes, 7–9
 - barriers, 9–10
- internal occlusion providers, 3
- iontophoresis, 367–373
 - cosmetic applications, 386–387
 - LFS (Low Frequency Sonophoresis), 373, 376–377
 - medical applications, 384–385
 - patches, 387–388
 - penetration enhancement, 366–367
 - principles, 382–384
 - safety, 388
- IR (infrared spectroscopic imaging), skin delivery, 49–57

L

leave-off products, 6

legal perspective, cosmetic and skin delivery, 462–464

antioxidants, 467

claims, 471–472

cosmeceuticals, 465–466

nanotechnology, 472–473

natural/organic products, 466

safety testing, 470–471

skin protectants, 467–468

test methods, 464–465

LFS (Low Frequency Sonophoresis)

iontophoresis, 373, 376–377

sonophoresis, 374–376

linoleic acid

molecular inclusion for oxidation protection, 357–358

physiological importance, 355–357

topical applications, 357

lipids

between corneocytes, 3

SLN[®] and NLC[®], 229–236

marketing SLN[®] cosmetic products, 244–245

pharmacokinetic and biopharmaceutic properties, 236–240

processing formulations, 240–244

liposomes, 13–14, 270–271. *See also* Ethosomes and Transfersomes

flexible, 271–278

nanoencapsulation, 318

liquid crystalline emulsions, 140, 142–155

clear gel-like emulsion crystals, 160–164

lemellar liquid gel, 164–168

molecular assemblies, 140–141

multilamellar emulsion of pseudo-stratum corneum lipids, 155–160

Low Frequency Sonophoresis. *See* LFS

M

marketing systems for skin delivery, 481–484

market size, 477–480

- necessity of skin care products, 480–481
- MEC (minimal effective concentration) process, 5
- medical drug delivery, biodegradable SLNs (solid-lipid nanoparticles), 331–334
- microencapsulation, 252–253, 314–315
 - chemical production, 316–317
 - physical production, 317
 - topical delivery systems, 338–341
 - polymeric entrapments, 341–350
- minimal effective concentration. *See* MEC
- moisturizers, 3

N

- nanoencapsulation
 - liposomes, 318
 - solid-lipid nanoparticles, 319–331
 - biodegradable, 331–334
- nanostructured lipid carriers. *See* NLC*
- NLC* (nanostructured lipid carriers), 229–236
 - marketing SLN* cosmetic products, 244–245
 - pharmacokinetic and biopharmaceutic properties, 236–240
 - processing formulations, 240–244

O

- OECD (Organization for Economic Cooperation and Development), skin penetration studies, 30, 31

P

- patches, 14–15
 - iontophoresis, 387–388
- percutaneous absorption, 30–32
 - CAID (cosmetic active ingredient delivery), 490–495
 - stratum corneum tape-stripping, 38–42
 - in vitro methodology, 32–38
 - versus* in vivo methodology, 42–43
- pharmaceutical drug delivery, 4
 - biodegradable SLNs (solid-lipid nanoparticles), 331–334
 - spectroscopy, acoustic attenuation, 186–202

Pheroids

- characteristics
 - functional, 293–300
 - structural, 284–293
 - cosmetic applications, 307–308
 - skin and Pheroid formulation compatibility, 304–307
 - therapeutic efficacy, 300–303
- polarity of stratum corneum, effects on dermal and transdermal cosmetic delivery, 110–122
- polyunsaturated fatty acids (vitamin F), 355
- pro-drug approach, delivery, 4–5

R

- rinse-off product delivery, 6, 403–405
- methods, 406–409
 - studies, 409–420
- RPI values, 97–98

S

- SCCNFP (Scientific Committee on Cosmetic Products and Non-Food, skin penetration studies, 30, 31
- shunt routes, skin penetration routes, 7–8
- skin delivery, 2
- legal perspective, 462–464
 - antioxidants, 467
 - claims, 471–472
 - cosmeceuticals, 465–466
 - nanotechnology, 472–473
 - natural/organic products, 466
 - safety testing, 470–471
 - skin protectants, 467–468
 - test methods, 464–465
 - marketing systems, 481–484
 - market size, 477–480
 - necessity of skin care products, 480–481
 - process overview, 6–7
 - relevance to cosmetic delivery, 7–8, 10
- skin electroporation
- combined with other methods, 398

- of cosmetics, 399
- dermaportation, 400
- of macromolecules, 396–397
- patient tolerance, 399
- of small- to moderately-sized molecules, 394–396
- ways to enhance, 392–394
- skin penetration
 - animal skin *versus* human skin, 26–27
 - delivery
 - dermal and transdermal, 16–18
 - follicular, 14
 - guidelines, 15–16
 - measuring, 23–27
 - percutaneous absorption, 30–42
 - routes, 7–9
- skin penetration from emulsions
 - emollient influence, 91–92
 - applied active ingredient doses, 103–104
 - “Formulating for Efficacy,” 98–106
 - skin delivery theory, 92–98
 - emulsifier influence, 126–138
- skin permeability
 - animal skin *versus* human skin, 26–27
 - measuring, 23–27
 - percutaneous absorption, 30–42
- SLN[®] (solid lipid nanoparticles), 229–236
 - marketing SLN[®] cosmetic products, 244–245
 - pharmacokinetic and biopharmaceutic properties, 236–240
 - processing formulations, 240–244
- SLNs (solid-lipid nanoparticles), 319–331
 - biodegradable, 331–334
 - liposomes, 318
- solid lipid nanoparticles. *See* SLN[®]
- solid-lipid nanoparticles. *See* SLNs
- sonophoresis
 - LFS (Low Frequency Sonophoresis), 373, 374–376
 - penetration enhancement, 366–367

- topical delivery, 373–375
- spectroscopy and skin delivery
 - acoustic attenuation, 186–202
 - IR (infrared spectroscopic imaging), 49–57
- stratum corneum, 11–12
 - cyclodextrins, 358–363
 - effects of changing polarity, on dermal and transdermal cosmetic delivery, 110–122
 - linoleic acid
 - molecular inclusion for oxidation protection, 357–358
 - physiological importance, 355–357
 - topical applications, 357
 - multilamellar emulsion of pseudo-stratum corneum lipids, 155–160
 - outside layer, 3
 - penetration routes (transepidermal transport), 7–8
 - barriers, 9–10
 - polyunsaturated fatty acids (vitamin F), 355
 - radioactivity distribution, 5
 - tape-stripping, percutaneous absorption, 38–42

T

- tape-stripping
 - versus* cyanoacrylate stripped skin, 449–451
 - stratum corneum, 38–42
- TDS (topical delivery systems). *See also* CAID
 - acoustic attenuation spectroscopy, 186–202
 - linoleic acid, 357
 - liposomes, flexible, 271–278
 - microencapsulation, 338–350
- therapeutic window, 6
- topical delivery systems. *See* TDS
- transcellular transepidermal penetration routes, 7–9
 - barriers, 9–10
- transdermal delivery, 4
 - avoiding, 10
 - changing polarity of stratum corneum effects on, 110–122
 - formulation type's influence, 77–80
 - measuring, 16–18